PROCEEDINGS OF SCIENCE

arXiv:1111.0336v1 [hep-lat] 1 Nov 2011

String tension at finite temperature Lattice QCD

Pedro Bicudo[*]
CFTP, Instituto Superior Técnico, Universidade Técnica de Lisboa
E-mail: `bicudo@ist.utl.pt`

Nuno Cardoso
CFTP, Instituto Superior Técnico, Universidade Técnica de Lisboa
E-mail: `nunocardoso@cftp.ist.utl.pt`

Orlando Oliveira
CFC, Universidade de Coimbra
E-mail: `orlando@teor.fis.uc.pt`

Paulo J. Silva
CFC, Universidade de Coimbra
E-mail: `psilva@teor.fis.uc.pt`

The critical curve of string tension as a function of the temperature is computed in SU(3) Lattice QCD. We present the results for the string tension utilizing a pair of Polyakov loop and antiloop, with two different techniques. We compare the colour averaged loop-antiloop which is gauge invariant but is only adequate to study the string tension, and the colour singlet loop-antiloop using the Landau gauge fixing which also enables to compute the coulomb part of the free energy.

The XXIX International Symposium on Lattice Field Theory - Lattice 2011
July 10-16, 2011
Squaw Valley, Lake Tahoe, California

---

[*] Speaker.

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike Licence.



Figure 1: $\sigma(T)$ Bielefeld data and emprirical curve

1. Introduction and Motivation

The confinement/deconfinement transition and chiral symmetry restoration are non-perturbative phenomena, elevant for heavy ion collisions and for quark models at finite temperature.

The nature of the confinement/deconfinement phase transition has been investigated in Lattice QCD using as order parameter the Polyakov Loop. For quenched QCD, the lattice data points towards first transition. However the single Polyakov loop is more adequate for detailed studies of temperatures $T > T_c$.

Here we study the string tension σ, adequate to study temperatures $T < T_c$. In the formulation of quark models [1], the *value of the string tension* σ determines the linear confining potential, with $\sigma \neq 0$ in the confined, low temperature phase and $\sigma = 0$ above the critical temperature T_c (deconfined phase). Further, *chiral symmetry* can also be related with σ [2].

Therefore, the string tension is also an order parameter for the deconfinement and chiral phase transition, which can be measured via two Polyakov loops.

The Bielefeld lattice QCD group measured the quark-antiquark free energy in a wide range of temperatures $T > 0.8T_c$ and also measured the critical curve $\sigma(T)$ [3, 4]. In Fig. 1 we depict the Bielefeld results, together with the empirical fit of Bicudo [5].

In order to understand the nature of the deconfinement phase transition, here we aim to measure $\sigma(T)$ also for temperatures smaller than $0.8T_c$.

To generate our configurations and to compute the Polyakov loops, we utilize GPUs of the NVIDIA FERMI generation [6, 7] and the TFlops cluster Milipeia [8].



Figure 2: Color average free energy with the link integration method



Figure 3: Color average free energy with the Landau gauge fixing



Figure 4: Color singlet free energy with the Landau gauge fixing

2. Framework

We extract the string tension form the static quark-antiquark free energy. The free energy is computed with a pair of Polyakov loop-antiloop. We explore two different color approaches to the Polyakov loop-antiloop, defined in Table 1.

The color average Polyakov loop-antiloop pair, leading to the color average free energy is gauge invariant, but at shorter distances the free energy may be contaminated by the color octet contribution.

On the other hand, the Singlet Free Energy is the one we aim at, however it is composed by the product of two non gauge invariant Wilson line, and in order to evaluate it we have to fix the gauge, choosing a particular gauge. Here we use the Landau gauge.

In this poster we apply both the gauge invariant calculations color average technique and we also utilize the Landau gauge technique.

We use the equation of the lattice spacing times the string tension at zero temperature, given

Polyakov Loop	$L(\vec{r})$	$= U_t(\vec{r},0)\,U_t(\vec{r},1)\cdots U_t(\vec{r},N_t-1)$
		$\propto \exp\left(-F_q/T\right)$
Singlet Correlator	$Re\left\{Tr\left[\langle L(\vec{0})L^\dagger(\vec{r})\rangle\right]\right\}$	$\propto \exp\left(-F_1(\vec{r},T)/T\right)$
Color Average Correlator	$\langle Re\left\{Tr\left[L(\vec{0})\right]\right\}\ Re\left\{Tr\left[L^\dagger(\vec{r})\rangle\right]\right\}$	$\propto \exp\left(-F_{ave}(\vec{r},T)/T\right)$
Singlet Free Energy	$F_1(\vec{r},T)$	
Octet Free Energy	$F_8(\vec{r},T)$	
Color Average Free Energy	$F_{ave}(\vec{r},T)$	$= \frac{1}{9}F_1 + \frac{8}{9}F_8$

Table 1: Color average and color singlet free energies.



Figure 5: $\sigma(T)$ for $T \in 0.48T_c - T_c$ extracted with the color average free energy

by

$$\left(a\sqrt{\sigma}\right)(g) = f(g^2)\left[1+0.2731\,\hat{a}(g)^2 - 0.01545\,\hat{a}(g)^4 + 0.01975\,\hat{a}(g)^6\right]/0.01364 \tag{2.1}$$

in Edwards et al. [9] valid in the region $5.6 \leq \beta \leq 6.5$,

$$\begin{aligned} \hat{a}(g) &= \frac{f(g^2)}{f(g^2(\beta=6.0))}, \\ f(g^2) &= \left(b_0 g^2\right)^{-\frac{b_1}{2b_0^2}} \exp\left(-\frac{1}{2b_0 g^2}\right), \\ b_0 &= \frac{11}{(4\pi)^2}, \qquad b_1 = \frac{102}{(4\pi)^4} \end{aligned} \tag{2.2}$$

Therefore, T/T_c is given by

$$\frac{T}{T_c} = \frac{\left(a\sqrt{\sigma}\right)(g(\beta_c))}{\left(a\sqrt{\sigma}\right)(g(\beta))} \tag{2.3}$$

where β_c was obtained in Lucini et al. [10].

The critical temperature [11, 12] T_c, in the quenched case is circa 270 MeV, for 2 flavour dynamical fermions 170 MeV and for 3 flavour dynamical fermions150 MeV. Here we study the critical curve $\sigma(T)$ in quenched SU(3) QCD.



Figure 6: $\sigma(T)$ for $T \in 0.48T_c - T_c$ extracted both with the color average free energy and with the color singlet free energy



Figure 7: Summary of ours and Bielefeld results for $\sigma(T)$ at $T \in 0.48T_c - T_c$

3. Results and conclusion

Our results are preliminary results only, aimed at exploring the string tension critical curve $\sigma(T)$ for finite temperatures below T_c, in particular for the so far unexplored $T < 0.8T_c$. We illustrate the different free energies in Figs 2, 3 and 4.

The noise leads to quite large error bars in the string tension points of Figs. 5, 6 and 7. We need to improve our signal to noise ratio, and to improve the precision of our temperatures, in order to improve the study of the critical curve.

Nevertheless, our preliminary results show that it is possible to study the critical curve of $\sigma(T)$ in a wider temperature range, and so far the empirical critical curve continues to show good agreement with the expected values.

Acknowledgments

This work was partly funded by the FCT contracts, POCI/FP/81933/2007, CERN/FP/83582/2008, PTDC/FIS/100968/2008, CERN/FP/109327/2009 and CERN/FP/116383/2010. Nuno Cardoso is also supported by FCT under the contract SFRH/BD/44416/2008. Paulo Silva is also supported by FCT under the contract SFRH/BPD/40998/2007.

References

[1] P. Bicudo, N. Cardoso and M. Cardoso, PoS B **ORMIO2011**, 062 (2011) [arXiv:1105.0063 [hep-ph]].

[2] P. Bicudo, arXiv:1109.6799 [hep-ph].

[3] O. Kaczmarek, F. Karsch, E. Laermann and M. Lutgemeier, Phys. Rev. D **62**, 034021 (2000) [arXiv:hep-lat/9908010].

[4] M. Doring, K. Huebner, O. Kaczmarek and F. Karsch, Phys. Rev. D **75**, 054504 (2007) [arXiv:hep-lat/0702009].

[5] P. Bicudo, Phys. Rev. D **82**, 034507 (2010) [arXiv:1003.0936 [hep-lat]].

[6] N. Cardoso and P. Bicudo, J. Comput. Phys. **230**, 3998 (2011) [arXiv:1010.4834 [hep-lat]].

[7] N. Cardoso, M. Cardoso and P. Bicudo, arXiv:hep-lat/1107.1355, (2000).

[8] Our computer facilities are described at the Portuguese Lattice QCD collaboration, and our CUDA codes are also available there. `http://nemea.ist.utl.pt/~ptqcd`.

[9] R. G. Edwards, U. M. Heller and T. R. Klassen, Nucl. Phys. B **517**, 377 (1998) [arXiv:hep-lat/9711003].

[10] B. Lucini, M. Teper and U. Wenger, JHEP **0401**, 061 (2004) [arXiv:hep-lat/0307017].

[11] Z. Fodor and S. D. Katz, JHEP **0203**, 014 (2002) [arXiv:hep-lat/0106002].

[12] R. Brower, P. Rossi and C. I. Tan, Nucl. Phys. B **190**, 699 (1981).